

Amino References:

Adams, GR, Haddad F. The relationships among IGF-1, DNA content, and protein accumulation during skeletal muscle hypertrophy. *J Appl Physiol* 1996; 81:2509-2516.

Alvestrand A, Hagenfeldt L, Merli M, Oureshi A, Eriksson LS. Influence of leucine infusion on intracellular amino acids in humans. *Eur J Clin Invest* 1990 Jun; 20(3):293-8.

Anthony JC, Anthony TG, Kimball SR, Vary TC, Jefferson LS Orally administered leucine stimulates protein synthesis in skeletal muscle of postabsorptive rats in association with increased eIF4F formation. *J Nutr* 2000; 130:139–145.

Anthony JC, Anthony TG, Layman DK Leucine supplementation enhances skeletal muscle recovery in rats following exercise. *J Nutr* 1999;129:1102–1106.

Anthony JC, Yoshizawa F, Anthony TG, Vary TC, Jefferson LS, Kimball SR. Leucine stimulates translation initiation in skeletal muscle of postabsorptive rats via a rapamycin-sensitive pathway. *J Nutr* 2000; 130:2413–2419.

Anthony JG, Fabian JR, Kimball SR, Jefferson LS. Identification of domains within the epsilon-subunit of the translation initiation factor eIF2B that are necessary for guanine nucleotide exchange activity and eIF2B holoprotein formation. *Biochim Biophys Acta* 2000 Jun 21; 1492(1):56-62.

Anthony JC, Lang CH Crozier SJ, Anthony TG, MacLean DA, Kimball SR, Jefferson LS. Contribution of insulin to the translational control of protein synthesis in skeletal muscle by leucine. *Am J Physiol Endocrinol Metab* 2002; 282:E1092-E1101.

Anthony, JC, Reiter AK, Anthony TG, Crozier SJ, Lang CH, MacLean DA, Kimball SR, Jefferson LS. Orally administered leucine enhances protein synthesis in skeletal muscle of diabetic rats in the absence of increases in 4E-BP1 or S6K1 phosphorylation. *Diabetes* 2002; 51:928-936.

Arfvidsson, B, Zachrisson H, Möller-Loswick AC, Hyltander A, Sandström R, Lundholm K. Effect of systemic hyperinsulinemia on amino acid flux across human legs in postabsorptive state. *Am J Physiol Endocrinol Metab* 1991; 260:E46-E52.

Augert, G, Monier S, Merchand-Brustel YL. Insulin affects only initiation and not elongation in protein synthesis in soleus muscles of lean and obese mice. *FEBS Lett* 1982; 147:211-214.

MetabolicDiet.com – Supplements

Baar, K, Esser K. Phosphorylation of p70s6k correlates with increased skeletal muscle mass following resistance exercise. *Am J Physiol Cell Physiol* 1999; 276:C120-C127.

Balage, M, Sinaud S, Prod'Homme M, Dardevet D, Vary TC, Kimball SR, Jefferson LS, Grizard J. Amino acids and insulin are both required to regulate assembly of the eIF4E · eIF4G complex in rat skeletal muscle. *Am J Physiol Endocrinol Metab* 2001; 281:E565-E574.

Balagopal, P, Schimke JC, Ades P, Adey D, Nair KS. Age effect on transcript levels and synthesis rate of muscle MHC and response to resistance exercise. *Am J Physiol Endocrinol Metab* 2001; 280:E203-E208.

Balon, TW, Zorzano A, Treadway JL, Goodman MN, Ruderman NB. Effect of insulin on protein synthesis and degradation in skeletal muscle after exercise. *Am J Physiol Endocrinol Metab* 1990; 258:E92-E97.

Bark, TH, McNurlan MA, Lang CH, Garlick PJ. Increased protein synthesis after acute IGF-1 or insulin infusion is localized to muscle in mice. *Am J Physiol Endocrinol Metab* 1998; 275:E118-E123.

Baron, AD, Steinberg H, Brechtel G, Johnson A. Skeletal muscle blood flow independently modulates insulin-mediated glucose uptake. *Am J Physiol Endocrinol Metab* 1994; 266:E248-E253.

Baron, AD. Hemodynamic action of insulin. *Am J Physiol Endocrinol Metab* 1994; 267: E187-E202.

Barrett EJ, Fryburg DA, Louard RJ, Gelfand RA, McNulty PH, Young LH Use of regional catheterization methods to study the regulation of muscle protein metabolism. In: Nair KS, ed. *Protein metabolism in diabetes mellitus*. London: Smith-Gordon 1992; 91–100

Barrett EJ, Revkin JH, Young LH, Zaret BL, Jacob R, Gelfand RA. An isotopic method for measurement of muscle protein synthesis and degradation in vivo. *Biochem J* 1987 Jul 1; 245(1):223-8.

Baumann, PQ, Stirewalt WS, O'Rourke BD, Howard D, Nair KS. Precursor pools of protein synthesis: a stable isotope study in a swine model. *Am J Physiol Endocrinol Metab* 1994; 267:E203-E209.

Bennet WM, Connacher AA, Jung RT, Stehle P, Rennie MJ. Effects of insulin and amino acids on leg protein turnover in IDDM patients. *Diabetes* 1991 Apr; 40(4):499-508.

MetabolicDiet.com – Supplements

Bennet WM, Connacher AA, Scrimgeour CM, Jung RT, Rennie MJ. Euglycemic hyperinsulinemia augments amino acid uptake by human leg tissues during hyperaminoacidemia. *Am J Physiol* 1990 Aug; 259(2 Pt 1):E185-94.

Bennet WM, Connacher AA, Scrimgeour CM, Rennie MJ. The effect of amino acid infusion on leg protein turnover assessed by L-[15N]phenylalanine and L-[1-13C] leucine exchange. *Eur J Clin Invest* 1990 Feb; 20(1):41-50.

Bennet WM, Connacher AA, Scrimgeour CM, Smith K, Rennie MJ. Increase in anterior tibialis muscle protein synthesis in healthy man during mixed amino acid infusion: studies of incorporation of [1-13C]leucine. *Clin Sci (Lond)* 1989 Apr; 76(4):447-54.

Bennet WM, Connacher AA, Smith K, Jung RT, Rennie MJ. Inability to stimulate skeletal muscle or whole body protein synthesis in type 1 (insulin-dependent) diabetic patients by insulin-plus-glucose during amino acid infusion: studies of incorporation and turnover of tracer L-[1-13C] leucine. *Diabetologia* 1990 Jan; 33(1):43-51.

Bergström, J. Factors causing catabolism in maintenance hemodialysis patients. *Miner Electrolyte Metab* 1992; 18:280-283.

Biolo G, Declan Fleming RY, Wolfe RR. Physiologic hyperinsulinemia stimulates protein synthesis and enhances transport of selected amino acids in human skeletal muscle. *J Clin Invest* 1995 Feb; 95(2):811-9.

Biolo G, Fleming RY, Maggi SP, Nguyen TT, Herndon DN, Wolfe RR. Inverse regulation of protein turnover and amino acid transport in skeletal muscle of hypercatabolic patients. *J Clin Endocrinol Metab* 2002 Jul; 87(7):3378-84.

Biolo G, Gastaldelli A, Zhang XJ, Wolfe RR. Protein synthesis and breakdown in skin and muscle: a leg model of amino acid kinetics. *Am J Physiol* 1994 Sep; 267(3 Pt 1):E467-74.

Biolo G, Iscra F, Bosutti A, Toigo G, Ciocchi B, Geatti O, Gullo A, Guarnieri G. Growth hormone decreases muscle glutamine production and stimulates protein synthesis in hypercatabolic patients. *Am J Physiol Endocrinol Metab* 2000 Aug; 279(2):E323-32.

Biolo G, Maggi SP, Williams BD, Tipton KD, Wolfe RR. Increased rates of muscle protein turnover and amino acid transport after resistance exercise in humans. *Am J Physiol* 1995 Mar; 268(3 Pt 1):E514-20.

Biolo G, Tipton KD, Klein S, Wolfe RR. An abundant supply of amino acids enhances the metabolic effect of exercise on muscle protein. *Am J Physiol* 1997 Jul; 273(1 Pt 1):E122-9.

MetabolicDiet.com – Supplements

Biolo G, Williams BD, Fleming RY, Wolfe RR. Insulin action on muscle protein kinetics and amino acid transport during recovery after resistance exercise. *Diabetes* 1999 May; 48(5):949-57.

Biolo, G, Fleming RYD, Maggi SP, Wolfe RR. Transmembrane transport and intracellular kinetics of amino acids in human skeletal muscle. *Am J Physiol Endocrinol Metab* 1995; 268:E75-E84.

Biolo, G, Fleming RYD, Wolfe RR. Physiologic hyperinsulinemia stimulates protein synthesis and enhances transport of selected amino acids in human skeletal muscle. *J Clin Invest* 1995; 95:811-819.

Biolo, G, Tipton KD, Klein S, Wolfe RR. An abundant supply of amino acids enhances the metabolic effect of exercise on muscle protein. *Am J Physiol Endocrinol Metab* 1997; 273:E122-E129.

Bodine, SC, Stitt TN, Gonzalez M, Kline WO, Stover GL, Bauerlein R, Zlotchenko E, Scrimgeour A, Lawrence JC, Jr, Glas DJ, Yancopoulos GD. AKT/mTOR pathway is a crucial regulator of skeletal muscle hypertrophy and can prevent muscle atrophy in vivo. *Nat Cell Biol* 2001; 3:1014-1019.

Bohe J, Low JFA, Wolfe RR, Rennie MJ Latency and duration of stimulation of human muscle protein synthesis during continuous infusion of amino acids. *J Physiol* 2001; 532:575–579.

Bonadonna RC, Saccomani MP, Cobelli C, DeFronzo RA. Effect of insulin on system A amino acid transport in human skeletal muscle. *J Clin Invest* 1993 Feb; 91(2):514-21.

Borah, MF, Schoenfeld PY, Gotch FA, Sargent JA, Wolfsen M, Humphreys MH. Nitrogen balance during intermittent dialysis therapy of uremia. *Kidney Int* 1978; 14:491-500.

Borsheim E, Tipton KD, Wolf SE, Wolfe RR. Essential amino acids and muscle protein recovery from resistance exercise. *Am J Physiol Endocrinol Metab* 2002 Oct; 283(4):E648-57.

Bradford M A rapid and sensitive method for the quantitation of microgram quantities of protein utilizing the principle of protein-dye binding. *Anal Biochem* 1976; 72:248–254.

Brand-Miller J. *The GI factor- the glucose revolution*. Hodder publishing 1998.

Brozinick, JT, Birnbaum MJ. Insulin, but not contraction activates AKT/PKB in isolated rat skeletal muscle. *J Biol Chem* 1998; 273:14679-14682.

MetabolicDiet.com – Supplements

Burgering BM, Coffey PJ Protein kinase B (c-Akt) in phosphatidylinositol-3-OH kinase signal transduction. *Nature* 1995; 376:599–602.

Burnett, PE, Barrow RK, Cohen NA, Snyder SH, Sabatini DM. RAFT1 phosphorylation of the translational regulators p70 S6 kinase and 4E-BP1. *Proc Natl Acad Sci USA* 1998; 95:1432-1437.

Buse MG. In vivo effects of branched chain amino acids on muscle protein synthesis in fasted rats. *Horm Metab Res* 1981; 13:502–505.

Buse MG, Weigand DA Studies concerning the specificity of the effect of leucine on the turnover of proteins in muscles of control and diabetic rats. *Biochim Biophys Acta* 1977; 475:81–89.

Bylund-Fellenius, A, Ojamaa KM, Flaim KE, Li JB, Wassner SJ, Jefferson LS. Protein synthesis versus energy state in contracting muscles of perfused rat hindlimb. *Am J Physiol Endocrinol Metab* 1984; 246:E297-E305.

Caiozzo, VJ, Haddad F, Baker MJ, Baldwin KM. Influence of mechanical loading on myosin heavy-chain protein and mRNA isoform expression. *J Appl Physiol* 1996; 80: 1503-1512.

Campbell, LE, Wang X, Proud CG. Nutrients differentially regulate multiple translation factors and their control by insulin. *Biochem J* 1999; 344: 433-441.

Caso G, Ford GC, Nair KS, Garlick PJ, McNurlan MA. Aminoacyl-tRNA enrichment after a flood of labeled phenylalanine: insulin effect on muscle protein synthesis. *Am J Physiol Endocrinol Metab* 2002 May; 282(5):E1029-38.

Caso G, Ford GC, Nair KS, Vosswinkel JA, Garlick PJ, McNurlan MA. Increased concentration of tracee affects estimates of muscle protein synthesis. *Am J Physiol Endocrinol Metab* 2001 Jun; 280(6):E937-46.

Castellino P, Luzi L, Simonson DC, Haymond M, DeFronzo RA. Effect of insulin and plasma amino acid concentrations on leucine metabolism in man. Role of substrate availability on estimates of whole body protein synthesis. *J Clin Invest* 1987 Dec; 80(6):1784-93.

Chesley, A, MacDougall JD, Tarnopolsky MA, Atkinson SA, Smith K. Changes in human muscle protein synthesis after resistance exercise. *J Appl Physiol* 1992; 73: 1383-1388.

Clark SE, Karn CA, Ahlrichs JA, Wang J, Leitch CA, Leichty EA, Denne SC. Acute changes in leucine and phenylalanine kinetics produced by parenteral nutrition in premature infants. *Pediatr Res* 1997 Apr; 41(4 Pt 1):568-74.

MetabolicDiet.com – Supplements

Coffer PJ, Jin J, Woodgett JR Protein kinase B (c-Akt): a multifunctional mediator of phosphatidylinositol 3-kinase activation. *Biochem J* 1998; 335:1–13.

Craddock BL, Price NT, Proud CG. Cloning and expression of cDNAs for the beta subunit of eukaryotic initiation factor-2B, the guanine nucleotide exchange factor for eukaryotic initiation factor-2. *Biochem J* 1995 Aug 1; 309(Pt 3):1009-14.

Dardevet D, Sornet C, Balage M, Grizard J. Stimulation of in vitro rat muscle protein synthesis by leucine decreases with age. *J Nutr* 2000 Nov; 130(11):2630-5.

Davis TA, Burrin DG, Fiorotto ML, Nguyen HV. Protein synthesis in skeletal muscle and jejunum is more responsive to feeding in 7-than in 26-day-old pigs. *Am J Physiol* 1996 May; 270(5 Pt 1):E802-9.

Davis, TA, Karl IE. Response of muscle protein turnover to insulin after acute exercise and training. *Biochem J* 1986; 240: 651-657.

Davis, TA, Nguyen HV, Suryawan A, Bush J, Jefferson LS, Kimball SR. Developmental changes in the feeding-induced stimulation of translation initiation in muscle of neonatal pigs. *Am J Physiol Endocrinol Metab* 2000; 279: E1226-E1234.

Di Pasquale MG. *Amino Acids and Proteins for the Athlete: The Anabolic Edge*. CRC Press 1997.

Dohm, GL, Tapscott EB, Barakat HA, Kasperek GJ. Measurement of in vivo protein synthesis in rats during an exercise bout. *Biochem Med* 1982; 27: 367-373.

Drenick, EJ, Swendseid ME, Bland WH, Tuttle SG. Prolonged starvation as a treatment of obesity. *JAMA* 1964; 187:100-105.

Dufner, A, Thomas G. Ribosomal S6 kinase signaling and the control of translation. *Exp Cell Res* 1999; 253: 100-109.

Fabian JR, Kimball SR, Jefferson LS. Reconstitution and purification of eukaryotic initiation factor 2B (eIF2B) expressed in Sf21 insect cells. *Protein Expr Purif* 1998 Jun; 13(1):16-22.

Fadden P, Haystead TAJ, Lawrence JC Identification of phosphorylation sites in the translational regulator, PHAS-I, that are controlled by insulin and rapamycin in rat adipocytes. *J Biol Chem* 1997; 272:10240–10247.

MetabolicDiet.com – Supplements

Fahien LA, MacDonald MJ. The succinate mechanism of insulin release. *Diabetes* 2002 Sep;51(9):2669-76.

Farrell PA, Hernandez JM, Fedele MJ, Vary TC, Kimball SR, Jefferson LS. Eukaryotic initiation factors and protein synthesis after resistance exercise in rats. *J Appl Physiol* 2000 Mar; 88(3):1036-42.

Farrell, PA, Fedele MJ, Hernandez J, Fluckey JD, Miller JL, III, Lang CH, Vary TC, Kimball SR, and Jefferson LS. Hypertrophy of skeletal muscle in diabetic rats in response to chronic resistance exercise. *J Appl Physiol* 1999; 87: 1083-1086.

Farrell, PA, Fedele MJ, Vary TC, Kimball SR, Lang CH, Jefferson LS. Regulation of protein synthesis after acute resistance exercise in diabetic rats. *Am J Physiol Endocrinol Metab* 1999; 276: E721-E727.

Farrell, PA, Hernandez JM, Fedele MJ, Vary TC, Kimball SR, Jefferson LS. Eukaryotic initiation factors and protein synthesis after resistance exercise in rats. *J Appl Physiol* 2000; 88: 1036-1042.

Fedele, MJ, Hernandez JM, Lang CH, Vary TC, Kimball SR, Jefferson LS, Farrell PA. Severe diabetes prohibits elevations in skeletal muscle protein synthesis following acute resistance exercise in rats. *J Appl Physiol* 88:102-108, 2000.

Fedele, MJ, Lang CH, Farrell PA. Immunization against IGF-1 prevents post-resistance exercise increases in protein synthesis in diabetic rats. *Am J Physiol Endocrinol Metab* 2001; 280:E877-E885.

Fedele, MJ, Vary TC, Farrell PA. IGF-I antibody prevents increases in protein synthesis in epitrochlearis muscles from refed, diabetic rats. *J Appl Physiol* 2001; 90:1166-1173.

Ferrando AA, Sheffield-Moore M, Paddon-Jones D, Wolfe RR, Urban RJ. Differential anabolic effects of testosterone and amino Acid feeding in older men. *J Clin Endocrinol Metab* 2003 Jan; 88(1):358-62.

Ferrando AA, Tipton KD, Doyle D, Phillips SM, Cortiella J, Wolfe RR. Testosterone injection stimulates net protein synthesis but not tissue amino acid transport. *Am J Physiol* 1998 Nov; 275(5 Pt 1):E864-71.

Ferrando AA, Williams BD, Stuart CA, Lane HW, Wolfe RR. Oral branched-chain amino acids decrease whole-body proteolysis. *JPEN J Parenter Enteral Nutr* 1995 Jan-Feb; 19(1):47-54.

Fluckey, JD, Vary TC, Jefferson LS, Evans WJ, Farrell PA. Insulin stimulation of protein synthesis in rat skeletal muscle following resistance exercise is maintained with advancing age. *J Gerontol Biol Sci* 1996; 51A:B323-B330.

MetabolicDiet.com – Supplements

Fluckey, JD, Vary TC, Jefferson LS, Farrell PA. Augmented insulin action on rates of protein synthesis after resistance exercise in rats. *Am J Physiol Endocrinol Metab* 1996; 270:E313-E319.

Fryburg DA, Barrett EJ, Louard RJ, Gelfand RA Effect of starvation on human muscle protein metabolism and its response to insulin. *Am J Physiol* 1990; 259:E477–E482

Fryburg DA, Jahn LA, Hill SA, Oliveras DM, Barrett EJ. Insulin and insulin-like growth factor-I enhance human skeletal muscle protein anabolism during hyperaminoacidemia by different mechanisms. *J Clin Invest* 1995 Oct; 96(4):1722-1729.

Fulks RM, Li JB, Goldberg AL Effects of insulin, glucose, amino acids on protein turnover in rat diaphragm. *J Biol Chem* 1975; 250:290–298.

Fumagalli, S, Thomas G. S6 phosphorylation and signal transduction. In: *Translational Control of Gene Expression*, edited by Mathews MB. Cold Spring Harbor, NY: Cold Spring Harbor Laboratory Press, 2000; p.695-717.

Gautsch, TA, Anthony JC, Kimball SR, Paul GL, Layman DK, Jefferson LS. Availability of eIF4E regulates skeletal muscle protein synthesis during recovery from exercise. *Am J Physiol Cell Physiol* 1998; 274:C409-C414.

Gelfand RA, Barrett EJ. Effect of physiologic hyperinsulinemia on skeletal muscle protein synthesis and breakdown in man. *J Clin Invest* 1987 Jul; 80(1):1-6.

Gingras, AC, Gygi SP, Raught B, Polakiewicz RD, Abraham RT, Hoekstra MF, Aebersold R, Sonenberg N. Regulation of 4E-BP1 phosphorylation: a novel two-step mechanism. *Genes Dev* 1999; 13:1422-1437.

Gingras, AC, Raught B, Gygi SP, Niedzwiecka A, Miron M, Burley SK, Polakiewicz RD, Wyslouch-Cieszynska A, Aebersold R, Sonenberg N. Hierarchical phosphorylation of the translation inhibitor 4E-BP1. *Genes Dev* 2001; 15:2852-2864.

Giordano M, Castellino P, DeFronzo RA Differential responsiveness of protein synthesis and degradation to amino acid availability in humans. *Diabetes* 1996; 45:393–399.

Giordano M, Castellino P, DeFronzo RA. Differential responsiveness of protein synthesis and degradation to amino acid availability in humans. *Diabetes*. 1996 Apr; 45(4):393-9.

MetabolicDiet.com – Supplements

Goldberg, AL. Influence of insulin and contractile activity on muscle size and protein balance. *Diabetes* 1979; 28, Suppl 1:18-24.

Goldberg, AL. Role of insulin in work-induced growth of skeletal muscle. *Endocrinology* 1968; 83:1071-1073.

Gomez E, Pavitt GD. Identification of domains and residues within the epsilon subunit of eukaryotic translation initiation factor 2B (eIF2Bepsilon) required for guanine nucleotide exchange reveals a novel activation function promoted by eIF2B complex formation. *Mol Cell Biol* 2000 Jun; 20(11):3965-76.

Greiwe JS, Kwon G, McDaniel ML, Semenkovich CF. Leucine and insulin activate p70 S6 kinase through different pathways in human skeletal muscle. *Am J Physiol* 2001; 281:E466–E471

Greiwe, JS, Kwon G, McDaniel ML, Semenkovich CF. Leucine and insulin activate p70 S6 kinase through different pathways in human skeletal muscle. *Am J Physiol Endocrinol Metab* 2001; 281:E466-E471.

Gross M, Rubino MS. Regulation of eukaryotic initiation factor-2B activity by polyamines and amino acid starvation in rabbit reticulocyte lysate. *J Biol Chem* 1989 Dec 25; 264(36):21879-84.

Guerrero-Ontivers, M.L. Wallimann, T. Creatine supplementation in Health and Disease. Effects of chronic creatine ingestion in vivo: Down regulation of the expression of creatine transporter isoforms in skeletal muscle. *Molecular and Cellular Biochemistry* 1998; 184:427-437.

Gutierrez A, Alvestrand A, Wahren J, Bergstrom J. Effect of in vivo contact between blood and dialysis membranes on protein catabolism in humans. *Kidney Int* 1990 Sep; 38(3):487-94.

Gutierrez A, Bergstrom J, Alvestrand A. Protein catabolism in sham-hemodialysis: the effect of different membranes. *Clin Nephrol* 1992 Jul; 38(1):20-9.

Hara K, Yonezawa K, Weng QP, Kozlowski MT, Belham C, Avruch J. Amino acid sufficiency and mTOR regulate P70 S6 kinase and eIF-4E BP1 through a common effector mechanism. *J Biol Chem* 1998; 273:14484–14494.

Hart, DW, Wolf SE, Chinkes DL, Gore DC, Mlcak RP, Beauford RB, Obeng MK, Lal S, Gold WF, Wolfe RR, Herndon DN. Determinants of skeletal muscle catabolism after severe burn. *Ann Surg* 2000; 232:455-465.

MetabolicDiet.com – Supplements

Hernandez, JM, Fedele MJ, Farrell PA. A time course evaluation of protein synthesis and glucose uptake and intracellular regulators after acute resistance exercise in rats. *J Appl Physiol* 2000; 88:1142-1149.

Hillier TA, Fryburg DA, Jahn LA, Barrett EJ. Extreme hyperinsulinemia unmasks insulin's effect to stimulate protein synthesis in the human forearm. *Am J Physiol* 1998 Jun; 274(6 Pt 1):E1067-74.

Iiboshi Y, Papst PJ, Kawasome H, Hosoi H, Abraham RT, Houghton PJ, Terada N. Amino acid-dependent control of p70S6K. Involvement of tRNA aminoacylation in the regulation. *J Biol Chem* 1999; 274:1092-1099.

Jacob R, Hu X, Niederstock D, Hasan S, McNulty PH, Sherwin RS, Young LH. IGF-I stimulation of muscle protein synthesis in the awake rat: permissive role of insulin and amino acids. *Am J Physiol* 1996 Jan; 270(1 Pt 1):E60-6.

Jefferson LS, Fabian JR, Kimball SR. Glycogen synthase kinase-3 is the predominant insulin-regulated eukaryotic initiation factor 2B kinase in skeletal muscle. *Int J Biochem Cell Biol* 1999 Jan; 31(1):191-200.

Jefferson LS, Kimball SR 2001 Amino acid regulation of gene expression. *J Nutr* 131:2460S-6S; discussion 2486S-2487S.

Jefferson LS, Li JB, Rannels SR. Regulation by insulin of amino acid release and protein turnover in the perfused rat hemikorpus. *J Biol Chem* 1977 Feb 25; 252(4):1476-83.

Jefferson, LS, Kimball SR. Amino acid regulation of gene expression. *J Nutr*; 2001; 131: 2460S-2466S.

Jefferson, LS, Rannels DE, Munger BL, Morgan HE. Insulin in the regulation of protein turnover in heart and skeletal muscle. *Fed Proc* 1974; 33:1098-1104.

Karinch AM, Kimball SR, Vary TC, Jefferson LS. Regulation of eukaryotic initiation factor-2B activity in muscle of diabetic rats. *Am J Physiol* 1993 Jan; 264(1 Pt 1):E101-8.

Karlstad MD, Buripakdi D, Carroll RC. Platelet-activating factor (PAF)-induced decreases in whole-body and skeletal muscle protein synthesis. *Shock* 2000 Oct; 14(4):490-8.

Kimball SR, Fabian JR, Pavitt GD, Hinnebusch AG, Jefferson LS. Regulation of guanine nucleotide exchange through phosphorylation of eukaryotic initiation factor eIF2 α . Role of the α - and δ -subunits of eIF2b. *J Biol Chem* 1998 May 22; 273(21):12841-5.

MetabolicDiet.com – Supplements

Kimball SR, Farrell PA, Jefferson LS. Invited Review: Role of insulin in translational control of protein synthesis in skeletal muscle by amino acids or exercise. *J Appl Physiol*. 2002 Sep; 93(3):1168-80.

Kimball SR, Heinzinger NK, Horetsky RL, Jefferson LS. Identification of interprotein interactions between the subunits of eukaryotic initiation factors eIF2 and eIF2B. *J Biol Chem* 1998 Jan 30; 273(5):3039-44.

Kimball SR, Horetsky RL, Jefferson LS Implication of eIF2B rather than eIF4E in the regulation of global protein synthesis by amino acids in L6 myoblasts. *J Biol Chem* 1998; 273(47):30945–53.

Kimball SR, Jefferson LS, Fadden P, Haystead TAJ, Lawrence JC Insulin and diabetes cause reciprocal changes in the association of eIF-4E and PHAS-I in rat skeletal muscle. *Am J Physiol* 1996; 270:C705–C709

Kimball, SR, Jefferson LS, Davis TA. Feeding stimulates protein synthesis in muscle and liver of neonatal pigs through an mTOR-dependent process. *Am J Physiol Endocrinol Metab* 2000; 279:E1080-E1087.

Kimball, SR, Vary TC, Jefferson LS. Regulation of protein synthesis by insulin. *Annu Rev Physiol* 1994; 56:321-348.

Kleijn M, Scheper GC, Voorma HO, Thomas AA Regulation of translation initiation factors by signal transduction. *Eur J Biochem* 1998; 253:531–544.

Kobayashi H, Borsheim E, Anthony TG, Traber DL, Badalamenti J, Kimball SR, Jefferson LS, Wolfe RR. Reduced amino acid availability inhibits muscle protein synthesis and decreases activity of initiation factor eIF2B. *Am J Physiol Endocrinol Metab* 2003 Mar; 284(3):E488-E498.

Kostyak JC, Kimball SR, Jefferson LS, Farrell PA. Severe diabetes inhibits resistance exercise-induced increase in eukaryotic initiation factor 2B activity. *J Appl Physiol* 2001 Jul; 91(1):79-84.

Kostyak, J, Kimball SR, Jefferson LS, Farrell PA. Severe diabetes inhibits the resistance exercise-induced increase in eukaryotic initiation factor 2B activity. *J Appl Physiol* 2001; 91:79-84.

Krishna VM, Janaki N, Ramaiah KV. Wheat germ initiation factor 2 (WG x eIF2) decreases the inhibition in protein synthesis and eIF2B activity of reticulocyte lysates mediated by eIF2alpha phosphorylation. *Arch Biochem Biophys* 1997 Oct 1; 346(1):28-36.

MetabolicDiet.com – Supplements

Lang CH, Frost RA, Jefferson LS, Kimball SR, Vary TC. Endotoxin-induced decrease in muscle protein synthesis is associated with changes in eIF2B, eIF4E, and IGF-I. *Am J Physiol Endocrinol Metab* 2000 Jun; 278(6):E1133-43.

Lang CH, Frost RA, Kumar V, Wu D, Vary TC. Impaired protein synthesis induced by acute alcohol intoxication is associated with changes in eIF4E in muscle and eIF2B in liver. *Alcohol Clin Exp Res* 2000 Mar; 24(3):322-31.

Lang CH, Frost RA, Nairn AC, MacLean DA, Vary TC. TNF-alpha impairs heart and skeletal muscle protein synthesis by altering translation initiation. *Am J Physiol Endocrinol Metab* 2002 Feb; 282(2):E336-47.

Laurent, GJ, Sparrow MP, Millward DJ. Turnover of muscle protein in the fowl. *Biochem J* 1978; 176:407-417.

Laurent, GL, Millward DJ. Protein turnover during skeletal muscle hypertrophy. *Fed Proc* 1980; 39:42-47.

Li JB, Jefferson LS Influence of amino acid availability on protein turnover in perfused skeletal muscle. *Biochim Biophys Acta* 1978; 544:351–359.

Lim, VS, Bier DM, Flanigan MJ, Sum-Ping ST. The effect of hemodialysis on protein metabolism. A leucine kinetic study. *J Clin Invest* 1993; 91:2429-2436.

Liu Z, Jahn LA, Long W, Fryburg DA, Wei L, Barrett EJ Branched chain amino acids activate messenger ribonucleic acid translation regulatory proteins in human skeletal muscle, and glucocorticoids blunt this action. *J Clin Endocrinol Metab* 2001; 86:2136–2143.

Lofberg E, Essen P, McNurlan M, Wernerman J, Garlick P, Anderstam B, Bergstrom J, Alvestrand A. Effect of hemodialysis on protein synthesis. *Clin Nephrol* 2000 Oct; 54(4):284-94.

Lofberg E, Gutierrez A, Wernerman J, Anderstam B, Mitch WE, Price SR, Bergstrom J, Alvestrand A. Effects of high doses of glucocorticoids on free amino acids, ribosomes and protein turnover in human muscle. *Eur J Clin Invest* 2002 May; 32(5):345-53.

Long W, Saffer L, Wei L, Barrett EJ Amino acids regulate skeletal muscle PHAS-I and p70 S6-kinase phosphorylation independently of insulin. *Am J Physiol* 2000; 279:E301–E306

Louard RJ, Barrett EJ, Gelfand RA Effect of infused branched-chain amino acids on muscle and whole body amino acid metabolism in man. *Clin Sci* 1990;79:457–466.

MetabolicDiet.com – Supplements

Louard RJ, Barrett EJ, Gelfand RA Overnight branched-chain amino acid infusion causes sustained suppression of muscle proteolysis. *Metab Clin Exp* 1995; 44:424–429.

Louard RJ, Barrett EJ, Gelfand RA. Effect of infused branched-chain amino acids on muscle and whole-body amino acid metabolism in man. *Clin Sci (Lond)* 1990 Nov; 79(5):457-66.

Louard RJ, Bhushan R, Gelfand RA, Barrett EJ, Sherwin RS Glucocorticoids antagonize insulin's antiproteolytic action on skeletal muscle. *J Clin Endocrinol Metab* 1994; 79:278–284.

MacDonald, MJ. Feasibility of a mitochondrial pyruvate malate shuttle in pancreatic islets. Further implication of cytosolic NADPH in insulin secretion. *J Biol Chem* 1995; 270: 20051-20058.

Markuns, JF, Wojtaszewski JFP, Goodyear LJ. Insulin and exercise decrease glycogen synthase kinase-3 activity by different mechanisms in rat skeletal muscle. *J Biol Chem* 1999; 274: 234896-249000.

McNulty PH, Young LH, Barrett EJ. Response of rat heart and skeletal muscle protein in vivo to insulin and amino acid infusion. *Am J Physiol* 1993 Jun; 264(6 Pt 1):E958-65.

Meek SE, Persson M, Ford GC, Nair KS. Differential regulation of amino acid exchange and protein dynamics across splanchnic and skeletal muscle beds by insulin in healthy human subjects. *Diabetes* 1998 Dec; 47(12):1824-35.

Milley JR. Effects of insulin on ovine fetal leucine kinetics and protein metabolism. *J Clin Invest* 1994 Apr; 93(4):1616-24.

Millward, DJ, Odedra B, Bates PC. The role of insulin, corticosterone and other factors in the acute recovery of muscle protein synthesis on refeeding food-deprived rats. *Biochem J* 1983; 216:583-587.

Mittendorfer B, Volpi E, Wolfe RR. Whole body and skeletal muscle glutamine metabolism in healthy subjects. *Am J Physiol Endocrinol Metab* 2001 Feb; 280(2):E323-33.

Moldawer LL, O'Keefe SJ, Bothe A Jr, Bistrian BR, Blackburn GL. In vivo demonstration of nitrogen-sparing mechanisms for glucose and amino acids in the injured rat. *Metabolism* 1980 Feb; 29(2):173-80.

Moller-Loswick AC, Zachrisson H, Hyltander A, Korner U, Matthews DE, Lundholm K. Insulin selectively attenuates breakdown of nonmyofibrillar proteins

MetabolicDiet.com – Supplements

in peripheral tissues of normal men. *Am J Physiol* 1994 Apr; 266(4 Pt 1):E645-52.

Moller-Loswick, AC, Zachrisson H, Hyltander A, Korner U, Matthews DE, Lundholm K. Insulin selectively attenuates breakdown of nonmyofibrillar proteins in peripheral tissues of normal men. *Am J Physiol Endocrinol Metab* 1994; 266:E645-E652.

Mortimore, GE, Poso AR, Kadowaki M, Wert JJ. Multiphasic control of hepatic protein degradation by regulatory amino acids: general features and hormonal modulation. *J Biol Chem* 1987; 262:16322-16327.

Mortimore, GE, Wert JJ, Miotto G, Venerando R, Kadowaki M. Leucine-specific binding of photoreactive Leu7-MAP to a high molecular weight protein on the plasma membrane of the isolated rat hepatocyte. *Biochem Biophys Res Commun* 1994; 203:200-208.

Nader, GA, Esser KA. Intracellular signaling specificity in skeletal muscle in response to different modes of exercise. *J Appl Physiol* 2001; 90:1936-1942 .

Nair KS, Ford GC, Ekberg K, Fernqvist-Forbes E, Wahren J. Protein dynamics in whole body and in splanchnic and leg tissues in type I diabetic patients. *J Clin Invest* 1995 Jun; 95(6):2926-37.

Narahara, HT, Holloszy JO. The actions of insulin, trypsin and electrical stimulation on amino acid transport in muscle. *J Biol Chem* 1974; 249:5435.

Nass R, Pezzoli SS, Chapman IM, Patrie J, Hintz RL, Hartman ML, Thorner MO. IGF-I does not affect the net increase in GH release in response to arginine. *Am J Physiol Endocrinol Metab*. 2002 Oct;283(4):E702-10.

Nave, BT, Ouwens DM, Withers DJ, Alessi DR, Shepherd PR. Mammalian target of rapamycin is a direct target for protein kinase B: identification of a convergence point for opposing effects of insulin and amino acid deficiency on protein translation. *Biochem J* 1999; 344:427-431.

O'Connor PM, Bush JA, Suryawan A, Nguyen HV, Davis TA. Insulin and amino acids independently stimulate skeletal muscle protein synthesis in neonatal pigs. *Am J Physiol Endocrinol Metab* 2003 Jan; 284(1):E110-9.

Ojamaa, KM, Kimball SR, Jefferson LS. Aminoacylation of initiator methionyl-tRNA_i under conditions inhibitory to initiation of protein synthesis. *Am J Physiol Endocrinol Metab* 1993; 264:E257-E263.

Pain, VM. Initiation of protein synthesis in eukaryotic cells. *Eur J Biochem* 1996;236:747-771.

MetabolicDiet.com – Supplements

Patti ME, Brambilla E, Luzi L, Landaker EJ, Kahn CR Bidirectional modulation of insulin action by amino acids. *J Clin Invest* 1998; 101:1519–1529.

Pause A, Belsham G, Gingras AC, Donze O, Lin TA, Lawrence JC, Sonenberg N Insulin-dependent stimulation of protein synthesis by phosphorylation of a regulator of 5'-cap function. *Nature* 1994; 371:762–767.

Pearson RB, Thomas G Regulation of p70S6K/p85S6K and its role in the cell cycle. *Prog Cell Cycle Res* 1995; 1:21–32.

Pham, PT, Heydrick SJ, Fox HL, Kimball SR, Jefferson LS, Lynch CJ. Assessment of cell signaling pathways in the regulation of mTOR by amino acids in rat adipocytes. *J Cell Biochem* 2000; 79:427-441.

Phillips, SM, Tipton KD, Aarsland A, Wolf SE, Wolfe RR. Mixed muscle protein synthesis and breakdown after resistance exercise in humans. *Am J Physiol Endocrinol Metab* 1997; 273:E99-E107.

Phillips, SM, Tipton KD, Ferrando AA, Wolfe RR. Resistance training reduces the acute exercise-induced increase in muscle protein turnover. *Am J Physiol Endocrinol Metab* 1999; 276:E118-E124.

Preedy, VR, Garlick PJ. The response of muscle protein synthesis to nutrient intake in postabsorptive rats: the role of insulin and amino acids. *Biosci Rep* 1986; 6:177-183.

Price NT, Kimball SR, Jefferson LS, Proud CG. Cloning of cDNA for the gamma-subunit of mammalian translation initiation factor 2B, the guanine nucleotide-exchange factor for eukaryotic initiation factor 2. *Biochem J* 1996 Sep 1; 318 (Pt 2):631-6.

Proud CG, Denton RM Molecular mechanisms for the control of translation by insulin. *Biochem J* 1997; 328:329–341.

Proud CG. p70 S6 kinase: an enigma with variations. *Trends Biochem Sci* 1996; 21:181–185.

Pullen, N, Dennis PB, Andjelkovic M, Dufner A, Kozma S, Hemmings BA, Thomas G. Phosphorylation and activation of p70s6k by PDK1. *Science* 1998; 279:707-710.

Rannels DE, Pegg AE, Rannels SR, Jefferson LS. Effect of starvation on initiation of protein synthesis in skeletal muscle and heart. *Am J Physiol* 1978 Aug; 235(2):E126-33.

MetabolicDiet.com – Supplements

Rasmussen BB, Tipton KD, Miller SL, Wolf SE, Wolfe RR. An oral essential amino acid-carbohydrate supplement enhances muscle protein anabolism after resistance exercise. *J Appl Physiol* 2000; 88(2):386-92.

Rasmussen BB, Wolfe RR, Volpi E. Oral and intravenously administered amino acids produce similar effects on muscle protein synthesis in the elderly. *J Nutr Health Aging* 2002; 6(6):358-62.

Ratheiser KM, Pesola GR, Campbell RG, Matthews DE. Epinephrine transiently increases amino acid disappearance to lower amino acid levels in humans. *JPEN J Parenter Enteral Nutr* 1999 Sep-Oct; 23(5):279-87.

Raught, B, Gingras AC, Sonenberg N. Regulation of ribosomal recruitment in eukaryotes. In: *Translational Control of Gene Expression*, edited by Mathews MB. Cold Spring Harbor, NY: Cold Spring Harbor Laboratory Press, 2000; p. 245-293.

Rennie MJ, Bohe J, Wolfe RR. Latency, duration and dose response relationships of amino acid effects on human muscle protein synthesis. *J Nutr* 2002 Oct;132(10):3225S-7S..

Rennie MJ. Control of muscle protein synthesis as a result of contractile activity and amino acid availability: implications for protein requirements. *Int J Sport Nutr Exerc Metab* 2001 Dec; 11 Suppl:S170-6.

Reynolds, TH, Bodine SC, Lawrence JC. Control of Ser2448 phosphorylation in the mammalian target of rapamycin by insulin and skeletal muscle load. *J Biol Chem* 2002; 277:17657-17662.

Rooyackers OE, Wagenmakers AJ. Tracer kinetics are of limited value to measure in vivo protein synthesis and degradation rates in muscle of anesthetized rats. *Metabolism*. 1996 Oct; 45(10):1279-83.

Sakurai Y, Aarsland A, Herndon DN, Chinkes DL, Pierre E, Nguyen TT, Patterson BW, Wolfe RR. Stimulation of muscle protein synthesis by long-term insulin infusion in severely burned patients. *Ann Surg*. 1995 Sep; 222(3):283-94; 294-7.

Sans MD, Kimball SR, Williams JA. Effect of CCK and intracellular calcium to regulate eIF2B and protein synthesis in rat pancreatic acinar cells. *Am J Physiol Gastrointest Liver Physiol* 2002 Feb; 282(2):G267-76.

Scheper GC, Mulder J, Kleijn M, Voorma HO, Thomas AA, van Wijk R. Inactivation of eIF2B and phosphorylation of PHAS-I in heat-shocked rat hepatoma cells. *J Biol Chem* 1997 Oct 24; 272(43):26850-6.

MetabolicDiet.com – Supplements

Scheper GC, Thomas AA, van Wijk R. Inactivation of eukaryotic initiation factor 2B in vitro by heat shock. *Biochem J.* 1998 Sep 1; 334 (Pt 2):463-7.

Schwartz J. Rapid modulation of protein synthesis in normal rats by specific neutralization and replacement of growth hormone. *Endocrinology* 1982 Dec; 111(6):2087-90.

Scorsone KA, Panniers R, Rowlands AG, Henshaw EC. Phosphorylation of eukaryotic initiation factor 2 during physiological stresses which affect protein synthesis. *J Biol Chem* 1987 Oct 25; 262(30):14538-43.

Scott PH, Brunn GJ, Kohn AD, Roth RA, Lawrence Jr JC 1998 Evidence of insulin-stimulated phosphorylation and activation of the mammalian target of rapamycin mediated by a protein kinase B signaling pathway. *Proc Natl Acad Sci USA* 95:7772–7777.

Sekulic, A, Hudson CC, Homme JL, Yin P, Otterness DM, Karnitz LM, Abraham RT. A direct linkage between the phosphoinositide 3-kinase-Akt signaling pathway and the mammalian target of rapamycin in mitogen-stimulated and transformed cells. *Cancer Res* 2000; 60:3504-3523.

Shah OJ, Anthony JC, Kimball SR, Jefferson LS. 4E-BP1 and S6K1: translational integration sites for nutritional and hormonal information in muscle. *Am J Physiol* 2000;279:E715–E729

Shah OJ, Antonetti DA, Kimball SR, Jefferson LS. Leucine, glutamine, and tyrosine reciprocally modulate the translation initiation factors eIF4F and eIF2B in perfused rat liver. *J Biol Chem* 1999 Dec 17; 274(51):36168-75.

Sheffield-Moore M, Urban RJ, Wolf SE, Jiang J, Catlin DH, Herndon DN, Wolfe RR, Ferrando AA. Short-term oxandrolone administration stimulates net muscle protein synthesis in young men. *J Clin Endocrinol Metab* 1999 Aug; 84(8):2705-11.

Sheffield-Moore M, Wolfe RR, Gore DC, Wolf SE, Ferrer DM, Ferrando AA. Combined effects of hyperaminoacidemia and oxandrolone on skeletal muscle protein synthesis. *Am J Physiol Endocrinol Metab* 2000 Feb; 278(2):E273-9.

Sherwood, DJ, Dufresne SD, Markuns JF, Cheatham B, Moller DE, Aronson D, Goodyear LJ. Differential regulation of MAP kinase, p70S6K, and Akt by contraction and insulin in rat skeletal muscle. *Am J Physiol Endocrinol Metab* 1999; 276:E870-E878.

Sinaud, S, Balage M, Bayle G, Dardevet D, Vary TC, Kimball SR, Jefferson LS, Grizard J. Diazoxide-induced insulin deficiency greatly reduced muscle protein

MetabolicDiet.com – Supplements

synthesis in rats: involvement of eIF4E. *Am J Physiol Endocrinol Metab* 1999; 276:E50-E61.

Singh LP, Wahba AJ. Regulation of protein synthesis in eukaryotic cells by the guanine nucleotide exchange factor and chain initiation factor 2. *SAAS Bull Biochem Biotechnol* 1996; 9:1-8.

Svanberg E, Moller-Loswick AC, Matthews DE, Korner U, Andersson M, Lundholm K. Effects of amino acids on synthesis and degradation of skeletal muscle proteins in humans. *Am J Physiol* 1996 Oct; 271(4 Pt 1):E718-24.

Svanberg, E, Jefferson LS, Lundholm K, Kimball SR. Postprandial stimulation of muscle protein synthesis is independent of changes in insulin. *Am J Physiol Endocrinol Metab* 1997; 272:E841-E847.

Svanberg, E, Zachrisson H, Ohlsson C, Iresjo BM, Lundholm KG. Role of insulin and IGF-I in activation of muscle protein synthesis after oral feeding. *Am J Physiol Endocrinol Metab* 1996; 270:E614-E620.

Tessari P, Barazzoni R, Zanetti M. Differences in estimates of forearm protein synthesis between leucine and phenylalanine tracers following unbalanced amino acid infusion. *Metabolism* 1999 Dec; 48(12):1564-9.

Tessari P, Garibotto G, Inchiostro S, Robaudo C, Saffioti S, Vettore M, Zanetti M, Russo R, Deferrari G. Kidney, splanchnic, and leg protein turnover in humans. Insight from leucine and phenylalanine kinetics. *J Clin Invest* 1996 Sep 15; 98(6):1481-92.

Tessari P, Zanetti M, Barazzoni R, Vettore M, Michielan F. Mechanisms of postprandial protein accretion in human skeletal muscle. Insight from leucine and phenylalanine forearm kinetics. *J Clin Invest* 1996 Sep 15; 98(6):1361-72.

Tipton KD, Ferrando AA, Phillips SM, Doyle D Jr, Wolfe RR. Postexercise net protein synthesis in human muscle from orally administered amino acids. *Am J Physiol* 1999 Apr; 276(4 Pt 1):E628-34.

Tipton KD, Wolfe RR. Exercise-induced changes in protein metabolism. *Acta Physiol Scand* 1998 Mar; 162(3):377-87.

Tipton, KD, Ferrando AA, Phillips SM, Doyle D, Wolfe RR. Postexercise net protein synthesis in human muscle from orally administered amino acids. *Am J Physiol Endocrinol Metab* 1999; 276:E628-E634.

Tipton, KD, Gurkin BE, Matin S, Wolfe RR. Nonessential amino acids are not necessary to stimulate net muscle protein synthesis in healthy volunteers. *J Nutr Biochem* 1999; 10:89-95.

MetabolicDiet.com – Supplements

Tipton, KD, Rasmussen PB, Miller SL, Wolf SE, Owens-Stovall SK, Petrini BE, Wolfe RR. Timing of amino acid-carbohydrate ingestion alters anabolic response of muscle to resistance exercise. *Am J Physiol Endocrinol Metab* 2001; 281:E197-E206.

Tremblay F, Marette A. Amino acid and insulin signaling via the mTOR/ p70 S6 kinase pathway. A negative feedback mechanism leading to insulin resistance in skeletal muscle cells. *J Biol Chem* 2001; 276:38052–38060.

Ueki K, Yamamoto-Honda R, Kaburagi Y, Yamauchi T, Tobe K, Burgering BMT, Coffey PJ, Komuro I, Akanuma Y, Yazaki Y, Kadowaki T. Potential role of protein kinase B in insulin-induced glucose transport, glycogen synthesis, and protein synthesis. *J Biol Chem* 1998; 273:5315–5322.

Vary TC, Jefferson LS, Kimball SR. Amino acid-induced stimulation of translation initiation in rat skeletal muscle. *Am J Physiol* 1999 Dec; 277(6 Pt 1):E1077-86.

Vary, TC, Jefferson LS, Kimball SR. Amino acid-induced stimulation of translation initiation in rat skeletal muscle. *Am J Physiol Endocrinol Metab* 1999; 277:E1077-E1086.

Vary, TC, Jefferson LS, Kimball SR. Insulin fails to stimulate muscle protein synthesis in sepsis despite unimpaired signaling to 4E-BP1 and S6K1. *Am J Physiol Endocrinol Metab* 2001; 281:E1045-E1053.

Volpi E, Ferrando AA, Yeckel CW, Tipton KD, Wolfe RR. Exogenous amino acids stimulate net muscle protein synthesis in the elderly. *J Clin Invest* 1998; 101(9):2000-7.

Volpi E, Mittendorfer B, Rasmussen BB, Wolfe RR. The response of muscle protein anabolism to combined hyperaminoacidemia and glucose-induced hyperinsulinemia is impaired in the elderly. *J Clin Endocrinol Metab* 2000 Dec; 85(12):4481-90.

Volpi E, Mittendorfer B, Wolf SE, Wolfe RR. Oral amino acids stimulate muscle protein anabolism in the elderly despite higher first-pass splanchnic extraction. *Am J Physiol* 1999 Sep; 277(3 Pt 1):E513-20.

Volpi E, Sheffield-Moore M, Rasmussen BB, Wolfe RR. Basal muscle amino acid kinetics and protein synthesis in healthy young and older men. *JAMA* 2001 Sep; 286(10):1206-12.

Wahren J, Felig P, Hagenfeldt L. Effect of protein ingestion on splanchnic and leg metabolism in normal man and in patients with diabetes mellitus. *J Clin Invest* 1976 ; 57:987–999.

MetabolicDiet.com – Supplements

Wang X, Campbell LE, Miller CM, Proud CG. Amino acid availability regulates p70 S6 kinase and multiple translation factors. *Biochem J* 1998;334:261–267.

Warner BW, Hasselgren PO, Hummel RP 3rd, James JH, Pedersen P, Fischer JE. Effect of catabolic hormone infusion on protein turnover and amino acid uptake in skeletal muscle. *Am J Surg* 1990 Mar; 159(3):295-300.

Watt PW, Corbett ME, Rennie MJ. Stimulation of protein synthesis in pig skeletal muscle by infusion of amino acids during constant insulin availability. *Am J Physiol* 1992 Sep; 263(3 Pt 1):E453-60.

Watt, PW, Goldspink G, Ward PS. Changes in fiber type composition in growing muscle as a result of dynamic exercise and static overload. *Muscle Nerve* 1984; 7:50-53.

Watt, PW, Kelly FJ, Goldspink DF, Goldspink G. Exercise-induced morphological and biochemical changes in skeletal muscles of the rat. *J Appl Physiol* 1982; 53:1144-1151.

Webb BL, Proud CG. Eukaryotic initiation factor 2B (eIF2B). *Int J Biochem Cell Biol.* 1997 Oct; 29(10):1127-31.

Welle, S, Bhatt K, Thornton CA. Stimulation of myofibrillar synthesis by exercise is mediated by more efficient translation of mRNA. *J Appl Physiol* 1999; 86:1220-1225.

Welsh GI, Miyamoto S, Price NT, Safer B, Proud CG. T-cell activation leads to rapid stimulation of translation initiation factor eIF2B and inactivation of glycogen synthase kinase-3. *J Biol Chem* 1996 May 10; 271(19):11410-3.

Welsh, GI, Miller CM, Loughlin AJ, Price NT, Proud CG. Regulation of eukaryotic initiation factor eIF2B: glycogen synthase kinase-3 phosphorylates a conserved serine which undergoes dephosphorylation in response to insulin. *FEBS Lett* 1998; 421:125-130.

Weng Q-P, Kozlowski M, Belham C, Zhang A, Comb M, Avruch J. Regulation of p70 S6 kinase by phosphorylation in vivo. *J Biol Chem* 1998; 273:16621–16629.

Williams, DD, Pavitt GD, Proud CG. Characterization of the initiation factor eIF2B and its regulation in *Drosophila melanogaster*. *J Biol Chem* 2001; 276:3733-3742.

Wolfe RR. Effects of amino acid intake on anabolic processes. *Can J Appl Physiol* 2001; 26 Suppl:S220-7.

MetabolicDiet.com – Supplements

Wolfe RR. Regulation of muscle protein by amino acids. *J Nutr* 2002 Oct; 132(10):3219S-24S..

Wolfe, RR, Miller SL. Amino acid availability controls muscle protein metabolism. *Diab Nutr Metab* 1999; 12:322-328.

Wolfe, RR, Volpi E. Insulin and protein metabolism. In: *Handbook of Physiology. The Endocrine Pancreas and Regulation of Metabolism*. Bethesda, MD: Am. Physiol. Soc 2001; sect. 7, vol. II, chapt.24, pages.735-757.

Wolfe, RR. *Radioactive and Stable Isotope Tracers in Biomedicine: Principles and Practice of Kinetic Analysis*. New York: Wiley-Liss, 1992.

Wong, TS, Booth FW. Protein metabolism in rat gastrocnemius muscle after stimulated chronic concentric exercise. *J Appl Physiol* 1990; 69:1709-1717.

Wong, TS, Booth FW. Protein metabolism in rat tibialis anterior muscle after stimulated chronic eccentric exercise. *J Appl Physiol* 1990; 69:1718-1724.

Wray-Cahen D, Nguyen HV, Burrin DG, Beckett PR, Fiorotto ML, Reeds PJ, Wester TJ, Davis TA. Response of skeletal muscle protein synthesis to insulin in suckling pigs decreases with development. *Am J Physiol*. 1998 Oct; 275(4 Pt 1):E602-9.

Yarasheski, KE, Zachwieja JJ, Bier DM. Acute effects of resistance exercise on muscle protein synthesis in young and elderly men and women. *Am J Physiol Endocrinol Metab* 1993; 265:E210-E214.

Yoshizawa F, Kimball SR, Jefferson LS Modulation of translation initiation in rat skeletal muscle and liver in response to food intake. *Biochem Biophys Res Commun* 1997; 240:825–831.

Yoshizawa F, Kimball SR, Vary TC, Jefferson LS Effect of dietary protein on translation initiation in rat skeletal muscle and liver. *Am J Physiol* 1998; 38:E814–E820

Yoshizawa, F, Kimball SR, Vary TC, Jefferson LS. Effect of dietary protein on translation initiation in rat skeletal muscle and liver. *Am J Physiol Endocrinol Metab* 1998; 275:E814-E820.

Yoshizawa, F, Nagasawa T, Nishizawa N, Funabiki R. Protein synthesis and degradation change rapidly in response to food intake in muscle of food-deprived mice. *J Nutr* 1997; 127:1156-1159.

MetabolicDiet.com – Supplements

Young LH, Dahl DM, Rauner D, Barrett EJ. Physiological hyperinsulinemia inhibits myocardial protein degradation in vivo in the canine heart. *Circ Res.* 1992 Aug; 71(2):393-400.

Young, VR, El-Khoury AE, Raguso CA, Forslund AH, Hambraeus L. Rates of urea production and hydrolysis and leucine oxidation change linearly over widely varying protein intakes in healthy adults. *J Nutr* 2000; 130:761-766.

Zorano A, Balon TW, Goodman MN, Ruderman NB. Insulin and exercise stimulate muscle alpha aminoisobutyric acid transport by a Na-K-ATPase pathway. *Biochem Biophys Res Commun* 1986; 134.

Zorano, A, Balon TW, Goodman MN, Garetto LP, Ruderman NB. Muscle - aminoisobutyric acid transport after exercise: enhanced stimulation by insulin. *Am J Physiol Endocrinol Metab* 1985; 248:E546-E552.